

Rational Solution of the KZ equation (example)

Andrey Tydnyuk

E-mail address: andrey.tydniouk@verizon.net
735 Crawford Ave. Brooklyn, NY 11223. USA.

Abstract

We investigate the Knizhnik-Zamolodchikov linear differential system. The coefficients of this system are rational functions. We prove that the solution of the KZ system is rational when k is equal to two and n is equal to three. While doing so, we found the coefficients of expansion in a neighborhood of a singular point.

Mathematics Subject Classification (2000). Primary 34M05, Secondary 34M55, 47B38.

Keywords: Symmetric group, linear differential system, rational solution.

Introduction

The Knizhnik-Zamolodchikov differential system has the form [1],[2]:

$$\frac{dW}{dz} = 2A(z)W, \quad (0.1)$$

where $A(z)$ and $W(z)$ are 3×3 matrices, $z_1 \neq z_2$. We suppose that $A(z)$ has the form

$$A(z) = \frac{P_1}{z - z_1} + \frac{P_2}{z - z_2}. \quad (0.2)$$

Here:

$$P_1 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (0.3)$$

$$P_2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \quad (0.4)$$

The matrices P_1 and P_2 are connected with the matrix representation of the symmetric group. In this paper, we consider the case when S_3 . We prove that in this case the solution of the Knizhnik-Zamolodchikov is rational. We find the coefficients of the Laurent expansion in the neighborhood of the point z_1 . We use the method of L. Sakhnovich [3].

1 MAIN NOTIONS

In a neighborhood of z_1 the matrix function $A(z)$ can be represented in the form:

$$A(z) = \frac{a_{-1}}{z - z_1} + a_0 + a_1(z - z_1) + \dots, \quad (1.1)$$

where

$$a_{-1} = P_1, \quad a_r = (-1)^r \frac{P_2}{(z_2 - z_1)^{r+1}}, \quad r \geq 0. \quad (1.2)$$

Proposition 1.1 (see[3])(necessary and sufficient condition) *If the matrix system*

$$[(q+1)I_3 - 2a_{-1}]b_{q+1} = 2 \sum_{j+\ell=q} a_j b_\ell, \quad -2 \leq q+1 \leq 2 \quad (1.3)$$

has a solution $b_{-2}, b_{-1}, b_0, b_1, b_2$ and $b_{-2} \neq 0$. Then system (1) has a solution

$$W(z) = \sum_{p \geq -2} b_p (z - z_1)^p, \quad b_{-2} \neq 0. \quad (1.4)$$

System (1.3) can be written in the following form

$$b_{-2} = I_3 - P_1 \quad (1.5)$$

$$b_{-1} = -2(I_3 + 2P_1)^{-1} a_0 b_{-2} \quad (1.6)$$

$$b_0 = -P_1(a_0 b_{-1} + a_1 b_{-2}) \quad (1.7)$$

$$b_1 = 2(I_3 - 2P_1)^{-1}(a_0 b_0 + a_1 b_{-1} + a_2 b_{-2}) \quad (1.8)$$

$$(I_3 - P_1)b_2 = (a_0 b_1 + a_1 b_0 + a_2 b_{-1} + a_3 b_{-2}) \quad (1.9)$$

Direct calculations show that:

$$b_{-2} = \begin{bmatrix} 1 & -1 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad (1.10)$$

$$b_{-1} = \frac{1}{-9(z_2 - z_1)} \begin{bmatrix} -12 & 12 & 0 \\ 6 & -6 & 0 \\ 6 & -6 & 0 \end{bmatrix}, \quad (1.11)$$

$$b_0 = \frac{1}{-9(z_2 - z_1)^2} \begin{bmatrix} 3 & -3 & 0 \\ -6 & 6 & 0 \\ 3 & -3 & 0 \end{bmatrix}, \quad (1.12)$$

$$b_1 = \frac{1}{-9(z_2 - z_1)^3} \begin{bmatrix} 6 & -6 & 0 \\ 6 & -6 & 0 \\ -12 & 12 & 0 \end{bmatrix}. \quad (1.13)$$

It follows from system (1.3) and the relations (1.10)-(1.13) that

$$(I_3 - P_1)b_2 = \frac{1}{-9(z_2 - z_1)^4} \begin{bmatrix} 1 & -1 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}. \quad (1.14)$$

It is easy to see that equation (1.14) has a solution. Using proposition 1 we obtain the statement.

Proposition 1.2 *Differential system (0.1) has a rational fundamental solution.*

References

1. Chervov A., Talalaev D., KZ equation, G-operators, quantum Drinfeld-Sokolov reduction and quantum Cayley-Hamilton identity, arXiv:hep-th/0607250, 2006.
2. Etingof P.I., Frenkel I.B., Kirillov A.A. jr., Lectures on Representation Theory and Knizhnik-Zamolodchikov Equations, Amer. Math. Society, 1998.
3. Sakhnovich L.A., Meromorphic Solutions of Linear Differential Systems, Painleve Type Functions. Preprint (to appear).